

Motion of a testing particle in gravitational field of a ring

Michal Demetrian*

Comenius University Mlynska Dolina M 105 Bratislava 4
842 48 Slovak Republic

February 2, 2008

Abstract

The planar motion of a testing body in the field of gravity of the massive ring is studied. The "perihelion" precession due to non-relativistic reasons is demonstrated on this simple example of planar motion in the radially symmetric (in the plane of motion) situation. The ring as the source of the gravitational field can serve as the toy model for quasi-planar sources of gravity appearing in the galactic dynamics, etc. The results are used to estimate the real perihelion shift of the planet Mercury due to gravitational quadrupole moment of the Sun.

1 The potential

Let us consider the thin (one dimensional) homogenous massive ring of radius R with total mass M . We consider the cartesian coordinates x, y, z such that the plane Oxy coincides with the plane defined by the ring itself, and the origin of our coordinates coincides with the center of the ring. Our first task is to compute the Newtonian potential $V = V(x, y)$ of the gravitational field of the ring in the plane Oxy . Because of the rotational invariance of the problem, the potential is the function of radial distance $r = \sqrt{x^2 + y^2}$ only. According to the Newton's gravitational law we have

$$\begin{aligned} V(r) &= -\frac{\kappa M}{2\pi R} \int_0^{2\pi} \frac{R d\psi}{[(R \cos(\psi) - x)^2 + (R \sin(\psi) - y)^2]^{1/2}} = \left| \begin{array}{l} x = r \cos(\phi) \\ y = r \sin(\phi) \end{array} \right| = \\ &= -\frac{\kappa M}{2\pi} \int_0^{2\pi} \frac{d\psi}{[R^2 + r^2 - 2Rr \cos(\psi - \phi)]^{1/2}} = -\frac{\kappa M}{2\pi} \int_0^{2\pi} \frac{d\psi}{[R^2 + r^2 - 2Rr \cos(\psi)]^{1/2}} = \\ &= -\frac{\kappa M}{2\pi} \frac{1}{\sqrt{R^2 + r^2}} \int_0^{2\pi} \frac{d\psi}{\left[1 - \frac{2Rr}{R^2 + r^2} \cos(\psi)\right]^{1/2}}. \end{aligned}$$

We see that the potential has the singularity at $r = R$ (at the ring). It holds:

$$X \equiv \frac{2Rr}{R^2 + r^2} = 2 \frac{\frac{r}{R}}{1 + \left(\frac{r}{R}\right)^2} \leq 1,$$

*email: demetrian@fmph.uniba.sk

and the sign of equality takes place only for $r = R$. We remind the definition of the complete elliptic integral of the first kind

$$K(x) = \int_0^{\pi/2} \frac{dy}{[1 - x \sin^2(y)]^{1/2}}, \quad 0 \leq x < 1,$$

with help of which we can express our potential as follows

$$\begin{aligned} \int_0^{2\pi} \frac{d\psi}{[1 - X \cos(\psi)]^{1/2}} &= 2 \int_0^{\pi} \frac{d\psi}{[1 - X \cos(\psi)]^{1/2}} = 2 \int_0^{\pi} \frac{d\psi}{[1 + X \cos(\psi)]^{1/2}} = \\ 2 \int_0^{\pi} \frac{d\psi}{[1 + X (\cos^2(\psi/2) - \sin^2(\psi/2))]^{1/2}} &= 2 \int_0^{\pi} \frac{d\psi}{[1 + X - 2X \sin^2(\psi/2)]^{1/2}} = \\ \frac{4}{\sqrt{1+X}} \int_0^{\pi/2} \frac{d\xi}{\left[1 - \frac{2X}{1+X} \sin^2(\xi)\right]^{1/2}} &= \frac{4}{\sqrt{1+X}} K\left(\frac{2X}{1+X}\right). \end{aligned}$$

Finally, we obtain the potential of the ring in the form

$$V(r) = -\frac{2\kappa M}{\pi R} \frac{1}{1 + \frac{r}{R}} K\left(\frac{4\frac{r}{R}}{(1 + \frac{r}{R})^2}\right). \quad (1)$$

Having the potential we can derive the gravitational force that acts upon a testing body of mass m :

$$f(r) \equiv mg(r) = -mV'(r) = -\frac{\kappa m M}{\pi R^2} \frac{(1 + \frac{r}{R}) E\left(\frac{4\frac{r}{R}}{(1 + \frac{r}{R})^2}\right) + (1 - \frac{r}{R}) K\left(\frac{4\frac{r}{R}}{(1 + \frac{r}{R})^2}\right)}{\frac{r}{R} \left[\left(\frac{r}{R}\right)^2 - 1\right]}, \quad (2)$$

where E is the complete elliptic integral defined for $0 \leq x \leq 1$ by

$$E(x) = \int_0^{\pi/2} \sqrt{1 - x \sin^2(y)} dy$$

and we have denoted by g the gravitational acceleration.




Figure 1: Plot of the potential (1) as a function of the fraction r/R with $2\kappa M/(\pi R) = 1$.

For the next purposes we will write down the approximative expressions of the potential in the following three cases:

- $r/R \gg 1$:

$$V(r) = -\frac{\kappa M}{r} \left\{ 1 + \frac{1}{4} \left(\frac{R}{r} \right)^2 + \frac{9}{64} \left(\frac{R}{r} \right)^4 + \mathcal{O} \left(\frac{R}{r} \right)^6 \right\} \quad (3)$$

- $r/R \ll 1$:

$$V(r) = -\frac{\kappa M}{R} \left\{ 1 + \frac{1}{4} \left(\frac{r}{R} \right)^2 + \frac{9}{64} \left(\frac{r}{R} \right)^4 + \mathcal{O} \left(\frac{r}{R} \right)^6 \right\} \quad (4)$$

- $r/R \approx 1$:

$$V(r) = -\frac{2\kappa M}{\pi R} \left\{ \left[\frac{\ln(2) - \gamma_E}{2} - \frac{1}{2} \frac{\Gamma'(1/2)}{\Gamma(1/2)} - \frac{1}{2} \ln \left| \frac{r}{R} - 1 \right| \right] + \left[\frac{1 + \gamma_E - \ln(2) + \Gamma'(1/2)/\Gamma(1/2)}{4} + \frac{1}{4} \ln \left| \frac{r}{R} - 1 \right| \right] \left(\frac{r}{R} - 1 \right) + \dots \right\}, \quad (5)$$

where $\gamma_E \approx 0.577216$ is the Euler's γ constant and Γ is the Euler's Gamma function.

Equation (3) tells us that at large distances from the ring the field is almost identical with the field of point mass M and the leading correction term (proportional to $(r/R)^2$) is the projection of the quadrupole field into the plane of the ring. Equation (4) shows that the center of the ring is an unstable equilibrium. If a testing particle is displaced a little bit from the center then it is forced to move towards the ring, and the force is proportional to the distance from the center (inverted linear harmonic potential). Finally, equation (5) shows the feature of the potential close the ring. Namely, there is the logarithmic divergency of the potential as $r \rightarrow R$.

2 Equations of motion and the conservation laws

An easy way how to obtain equations of motion is to write down the Lagrange function (the Lagrangian) of the testing particle with the mass m in polar coordinates (r, ϕ) (we will consider the motion in the plane of the ring only, as we have already mentioned above):

$$L = \frac{1}{2} m (\dot{r}^2 + r^2 \dot{\phi}^2) - mV(r), \quad (6)$$

where the dot stands for the time derivative, and derive the equations of motion as the Euler - Lagrange equations

$$\ddot{r} + V'(r) = 0, \quad r^2 \ddot{\phi} + 2r\dot{r}\dot{\phi} = 0. \quad (7)$$

The order of this system of two differential equations can be reduced using the conservation laws. First of all, the energy E is conserved:

$$\frac{1}{2} m (\dot{r}^2 + r^2 \dot{\phi}^2) + mV(r) = E. \quad (8)$$

The angular momentum vector must have the form $(0, 0, \mathcal{L})$. Since L does not depend on ϕ , we have the following expression for the z - component of the angular momentum:

$$mr^2 \dot{\phi} = \mathcal{L}. \quad (9)$$

The previous two equations can be combined into one expressing the radial velocity $\dot{r}$ as the function of the radial position r and the conserved quantities E and $\mathcal{L}$:

$$|\dot{r}| = \left[\frac{2E}{m} - 2V(r) - \frac{\mathcal{L}^2}{m^2 r^2} \right]^{1/2}. \quad (10)$$

The inequality

$$\frac{2E}{m} - 2V(r) - \frac{\mathcal{L}^2}{m^2 r^2} \geq 0$$

defines, for given values of E and $\mathcal{L}$, classically accessible region, i.e. the minimal r_m and the maximal r_M value of radial coordinate of considered particle.

3 Radial motion (free fall)

The radial motion or the free fall is characterized by the condition $\mathcal{L} = 0$. We will consider the situation when the initial (radial) velocity equals zero. This means that $E/m = V(r_0)$, where r_0 is the initial position, and equation (10) becomes

$$|\dot{r}| = \sqrt{2}[V(r_0) - V(r)]^{1/2}.$$

Our task is to compute the time of the fall to the ring T . First, we will consider the case when $r_0 > R$. Thus,

$$\dot{r} = -\sqrt{2}[V(r_0) - V(r)]^{1/2} \Rightarrow T(r_0) = \frac{1}{\sqrt{2}} \int_R^{r_0} \frac{dr}{[V(r_0) - V(r)]}.$$

Explicitly,

$$T(r_0) = \frac{1}{2} \left[\frac{\pi R^3 (1 + \frac{r_0}{R})}{\kappa M K \left(\frac{4r_0/R}{(1 + \frac{r_0}{R})^2} \right)} \right]^{1/2} \int_1^{r_0/R} \frac{dX}{\left[\frac{1 + \frac{r_0}{R}}{1 + X} \frac{K \left(\frac{4X}{(1 + X)^2} \right)}{K \left(\frac{4r_0/R}{(1 + r_0/R)^2} \right)} - 1 \right]^{1/2}}. \quad (11)$$

If the initial position of the particle is inside the ring ($r_0 < R$) then

$$\dot{r} = +\sqrt{2}[V(r_0) - V(r)]^{1/2} \Rightarrow T(r_0) = \frac{1}{\sqrt{2}} \int_R^{r_0} \frac{dr}{[V(r_0) - V(r)]},$$

thus (11) holds also in this case. The free fall at the ring from outside can be compared with the radial fall at the point mass M placed in the center of the ring. The time of duration of such a fall is given by

$$T_p(r_0) = \int_0^{r_0} \frac{dr}{\left[2\kappa M \left(\frac{1}{r} - \frac{1}{r_0} \right) \right]^{1/2}} = \left[\frac{r_0^3}{2\kappa M} \right]^{1/2} \int_0^1 \sqrt{\frac{\xi}{1 - \xi}} d\xi = \frac{\pi}{2} \left[\frac{r_0^3}{2\kappa M} \right]^{1/2}. \quad (12)$$

The quantities T and T_p are compared in the fig. 2.

4 Circular orbits

For a testing body on a circular orbit the centripetal force f_c must be equal to the gravitational force (2). This means that

$$f_c = ma_c = -m\dot{\phi}^2 r = -\frac{\mathcal{L}^2}{mr^3} \Rightarrow -\frac{\mathcal{L}^2}{mr^3} = g.$$

The last equation can obviously be fulfilled for any $r > R$ but, of course, cannot be fulfilled for any $r < R$, i.e. the testing particle can orbit the ring along any circle from outside.




Figure 2: Left graph shows the dependence of the free fall duration T on initial distance r_0/R according (11), where we have chosen for definiteness: $\kappa M = 1$ and $R = 1$. Right graph shows the fraction T/T_p with the parameters chosen in the same way as in the left graph. We see that the quantity T approaches relatively T_p as r_0/R grows. More detailed analysis would show that the difference $T_p - T$ approaches a positive constant of the order of unity as r_0/R runs to infinity.

5 Small deformations of circular orbits - "perihelion" shift

In this section we will consider the close-to-circular trajectory of a testing particle sufficiently distanced from the ring. This means that the potential of the ring can be approximated by the first two terms of (3):

$$V \approx -\frac{\kappa M}{r} \left(1 + \frac{1}{4} \frac{R^2}{r^2} \right) \equiv -\frac{\kappa M}{r} + \frac{\gamma}{r^3}, \quad \gamma = -\frac{1}{4} \kappa M R^2. \quad (13)$$

It is well-known that the closed non-circular trajectories of testing particle in a central field exist only in two cases: $V \sim r^2$ and $V \sim -1/r$, see eg. [1] or detailed proof in [2]. Therefore our close-to-circular orbit will not be closed, however for a small deviation from the circle it can be regarded as slowly rotating ellipse. We are interested in the angle connecting the two nearest positions of the particle and the ring (the point of closest approach of a planet (from our solar system, of course) and the Sun is called perihelion) and the center of our coordinate system. Combining the eqs. (9) and (10) we easily can derive that the change of the polar angle $\Delta\phi$ between the two forthcoming closest approaches of the particle with respect to the ring is given by

$$\begin{aligned} \Delta\phi &= 2 \int_{r_m}^{r_M} \frac{\frac{\mathcal{L}}{m} \frac{1}{r^2} dr}{\sqrt{\frac{2E}{m} - 2V - \frac{\mathcal{L}^2}{m^2 r^2}}} = 2\mathcal{L} \int_{r_m}^{r_M} \frac{\frac{1}{r^2} dr}{\sqrt{2m(E - V) - \frac{\mathcal{L}^2}{r^2}}} = \\ &- 2 \frac{\partial}{\partial \mathcal{L}} \int_{r_m}^{r_M} \sqrt{2m(E - V) - \frac{\mathcal{L}^2}{r^2}} dr, \end{aligned} \quad (14)$$

where r_m and r_M are the minimal and the maximal values of the radius vector of the particle, respectively. The angle difference $\Delta\phi$ can be written as: $\Delta\phi = 2\pi + \delta\phi$, where the difference 2π comes from the Newton potential and additional term $\delta\phi$ is called perihelion shift in the mechanics of planetary motion around the Sun or the periastron shift in the mechanics of binary stars. An illustration of the "perihelion shift" in our case is given in fig. 3.




Figure 3: Plot of the trajectory of a testing particle in the field of massive ring. The radius of the ring is $R = 1$ and E and $\mathcal{L}$ were chosen so that the related Keplerian trajectory would be described by $r_m = 3$ and $r_M = 15$. The body of the ring is depicted as the central circle. The particle orbits the ring anti-clockwise, we see, from the left figure, that the "perihelion" moves in the same direction.

Using (13) and the Taylor expansion of the subintegral expression in (14) we obtain

$$\delta\phi = \frac{\partial}{\partial \mathcal{L}} \frac{2m}{\mathcal{L}} \int_0^\pi r^2 \frac{m\gamma}{r^3} d\phi = 2m^2\gamma \frac{\partial}{\partial \mathcal{L}} \frac{1}{\mathcal{L}} \int_0^\pi \frac{1}{r(\phi)} d\phi, \quad (15)$$

where $r = r(\phi)$ is the elliptic trajectory in the Newtonian field of a point mass. For this trajectory we have

$$r = \frac{p}{1 + \epsilon \cos(\phi)} \quad p = \frac{\mathcal{L}^2}{\kappa m^2 M} \quad \epsilon = \left[1 + \frac{2E\mathcal{L}^2}{\kappa^2 m^3 M^2} \right]^{1/2},$$

where ϵ is the eccentricity and p is the parameter of the ellipse with the semi-axis

$$a = \frac{p}{1 - \epsilon^2} \quad b = \frac{p}{\sqrt{1 - \epsilon^2}},$$

and

$$r_m = \frac{p}{1 + \epsilon} \quad r_M = \frac{p}{1 - \epsilon}.$$

Finally, we have

$$\delta\phi = 2\pi m^2\gamma \frac{\partial}{\partial \mathcal{L}} \frac{1}{\mathcal{L}p} = -6\pi m^2\gamma \frac{1}{\mathcal{L}^2 p} = -6\pi \frac{(\kappa m M)m^3\gamma}{\mathcal{L}^4} = \frac{3}{2}\pi \frac{\kappa^2 m^4 M^2 R^2}{\mathcal{L}^4}, \quad (16)$$

or using the relation

$$a(1 - \epsilon^2) = \frac{\mathcal{L}^2}{\kappa m^2 M}$$

we can express the perihelion shift (16) with help of geometrical parameters of considered perturbed ellipse:

$$\delta\phi = \frac{3}{2}\pi \frac{R^2}{a^2(1 - \epsilon^2)^2}.$$

We have studied in details the motion of a testing particle in the gravitational field of the massive planar ring. As the main aim, we have determined the shift of the perihelion of the testing particle with non-circular orbit with sufficiently large radius. For $r/R \gg 1$ the gravitational field of the ring is practically identical with the field of central monopole of the mass M and the quadrupole field, see (3). The formula (16) describes, in fact, the perihelion shift for the particle in the planar motion due to quadrupole perturbation of Newton potential. Hence, it can be used to estimate the perihelion shift of the planet Mercury due to Solar quadrupole moment, considering the approximation that the trajectory of Mercury is placed in the plane of the solar equator. Following the results about the multipole expansion of the gravitational field of the Sun we have

$$V_{\odot} = -\frac{\kappa M_{\odot}}{r} \left\{ 1 - J_2 \frac{R_{\odot}^2}{r^2} P_2(\cos(\vartheta)) + \mathcal{O}(R_{\odot}^3/r^3) \right\},$$

where $M_{\odot}$ is the mass of the Sun, $R_{\odot}$ is the mean radius of the Sun, $P_2(z) = -1/2 + 3/2z^2$ is the Legendre polynomial that depends on the azimuthal angle ϑ and $J_2 = (2.18 \pm 0.06) \times 10^{-7}$ is the dimensionless quadrupole moment of the Sun - for a long time the value of solar gravitational quadrupole was very uncertain, this value is from [3]. In the equator of the Sun plane we have $\vartheta = \pi/2$ and therefore in the mentioned plane the potential has the form

$$V_{\odot} = -\frac{\kappa M_{\odot}}{r} \left\{ 1 + \frac{J_2}{2} \frac{R_{\odot}^2}{r^2} + \mathcal{O}(R_{\odot}^3/r^3) \right\}.$$

So, we can use the formula (16) to compute the Mercury's perihelion shift due to the Sun's quadrupole moment identifying the constant γ as follows:

$$\gamma = -\frac{1}{2} J_2 \kappa M_{\odot} R_{\odot}^2. \quad (17)$$

Finally, taking into account the Mercury's parameters: the mass $m \approx 3.302 \times 10^{23} kg$, distance from the Sun in the perihelion $r_m \approx 4.6 \times 10^7 km$, orbital velocity in the perihelion $v_M \approx 59 km/s$ ($\mathcal{L} = m r_m v_M$, orbital period is about 88 days), and the mass of the Sun: $M_{\odot} \approx 2 \times 10^{30} kg$ and its radius $R_{\odot} \approx 7 \times 10^5 km$, we have

$$\delta\phi = 3\pi J_2 \frac{\kappa^2 m^4 M_{\odot}^2 R_{\odot}^2}{\mathcal{L}^2} = 3\pi J_2 \frac{\kappa^2 M_{\odot}^2 R_{\odot}^2}{r_m^4 v_M^4} \approx 3.3 \times 10^{-10}. \quad (18)$$

This means that during the period of 100 years, the perihelion of the Mercury is shifted in the direction of planetary rotation around the Sun in approximately 0.28 arcsec. This value can be compared with the well-known value of the relativistic shift of the perihelion that is about 35 arcsec per a century. Our value (18) is quite less that is in accordance with accepted fact that the Solar quadrupole does not effect essentially the Mercury's orbit in the time interval about 100 years.

References

- [1] L.D. Landau and E.M. Lifshitz, *Course of Theoretical Physics: Mechanics*, Butterworth-Heinemann, (1976).
- [2] V.I. Arnold, *Mathematical Methods of Classical Mechanics*, Springer, sec. ed., (1977).
- [3] M. Stix, *The Sun*, Springer, sec. ed., (2002).